

BPGM Rabbit mAb [9KMU]

Cat NO. :A11869

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	н	P07738	30 kda	Rabbit	IgG	100ul,200ul

Applications detail:	Application	Dilution			
	WB	1:1000-2000			
	The optimal dilutions should be determined by the end user				
		_			
Conjugate:					

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human BPGM.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Expressed in red blood cells. Expressed in non-erythroid cells of the placenta, present in the syncytiotrophoblast layer of the placental villi at the feto-maternal interface (at protein level)..

Subcellular location:

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey VIr: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Plays a major role in regulating hemoglobin oxygen affinity by controlling the levels of its allosteric effector 2,3-bisphosphoglycerate (2,3-BPG). Also exhibits mutase (EC 5.4.2.11) activity..

Validation Data:

BPGM Rabbit mAb [9KMU] Images



Western blot (SDS PAGE) analysis of extracts from Human placenta tissue lyaste.using BPGM Rabbit mAb [9KMU] at dilution of 1:1000 incubated at 4° C over night

View more information on http://naturebios.com