

ADH4 Mouse mAb[CA4T]

Cat NO. :A12526

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	н	P08319	40kda	Mouse	IgG	100ul,200ul

Applications detail:

Application

WB

1:1000-2000

The optimal dilutions should be determined by the end user

\sim	•		gate:			
1''	าทแ	100	to.			
\mathbf{v}	JIII	Jua	LE:			

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human ADH4.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Subcellular location:

Cytoplasm.

Function:

Catalyzes the NAD-dependent oxidation of either all-trans-retinol or 9-cis-retinol (PubMed:17279314). Also oxidizes long chain omega-hydroxy fatty acids, such as 20-HETE, producing both the intermediate aldehyde, 20-oxoarachidonate and the end product, a dicarboxylic acid, (5Z,8Z,11Z,14Z)-eicosatetraenedioate (PubMed:16081420). Also catalyzes the reduction of benzoquinones (PubMed:10514444)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

ADH4 Mouse mAb[CA4T] Images



Western blot (SDS PAGE) analysis of extracts from HepG2 cells.Using ADH4 Mouse mAb IgG [CA4T] at dilution of 1:1000 incubated at $4^{\circ}\mathrm{C}$ over night.

View more information on http://naturebios.com