

ME3 Rabbit mAb [4V0T]

Cat NO. :A14514

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	H,M,R	Q16798	67 kDa	Rabbit	IgG	100ul,200ul

Applications detail: Application Dilution

WB 1:1000-2000

IHC 1:100,

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human ME3.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Expressed predominantly in organs with a low-division rate.

Subcellular location:

Mitochondrion matrix.

Function:

Catalyzes the oxidative decarboxylation of (S)-malate to pyruvate using NADP(+) as a cofactor (PubMed:7818469). Can also reverse the decarboxylation reaction, but only with significantly lower efficiency (PubMed:7818469)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

ME3 Rabbit mAb [4V0T] Images



Western blot (SDS PAGE) analysis of extracts from $\;$ Fetal brain tissue lyaste.using ME3 Rabbit mAb [4V0T] $\;$ at dilution of 1:1000 incubated at $4\,^\circ\!\mathrm{C}$ over night

View more information on http://naturebios.com