

# P-Cadherin Mouse mAb [C620]

Cat NO. :A21903

#### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	н	P22223	120 kDa	Mouse	IgG	100ul,200ul

Applications detail:

Application

WB

1:1000-2000

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

**Purification**:

Protein A purification

#### Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of Human P-Cadherin

#### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$ 

### Tissue specificity:

Expressed in some normal epithelial tissues and in some carcinoma cell lines..

### Subcellular location:

Cell membrane, Single-pass type I membrane protein.

#### Function:

Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells, cadherins may thus contribute to the sorting of heterogeneous cell types.


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



## **Validation Data:**

## P-Cadherin Mouse mAb [C620] Images



Western blot (SDS PAGE) analysis of extracts from A431 cells. Using P-Cadherin mouse mAb[C620] at dilution of 1:1000 incubated at  $4^{\circ}$ C over night.

View more information on http://naturebios.com