

# MYH10 Mouse mAb[22D9]

Cat NO. :A21908

### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	H,M,R	P35580	230kda	Mouse	IgG	100ul,200ul

Applications detail:

Application Dilution
WB 1:1000-2000
IHC 1:100
ICC/IF 1:100
The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

**Purification**:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human MYH10.

### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

## Tissue specificity:

Isoform 1 is expressed in cerebellum and spinal chord. Isoform 2 is expressed in cerebrum and retina. Isoform 3 is expressed in the cerebrum and to a much lower extent in cerebellum..

## Subcellular location:

Cell projection, lamellipodium.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Cellular myosin that appears to play a role in cytokinesis, cell shape, and specialized functions such as secretion and capping. Involved with LARP6 in the stabilization of type I collagen mRNAs for CO1A1 and CO1A2. During cell spreading, plays an important role in cytoskeleton reorganization, focal contacts formation (in the central part but not the margins of spreading cells), and lamellipodial extension, this function is mechanically antagonized by MYH9..

## **Validation Data:**

### MYH10 Mouse mAb[22D9] Images



View more information on http://naturebios.com