TNNI2 Rabbit mAb [4H42]

Cat NO. :A27454

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	H,M,R	P48788	21 kda	Rabbit	lgG	50ul,100ul,200ul

Applications detail:

Application Dilution WB 1:1000-2000 The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human TNNI2.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Subcellular location:

Function:

Troponin I is the inhibitory subunit of troponin, the thin filament regulatory complex which confers calciumsensitivity to striated muscle actomyosin ATPase activity.


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Validation Data:

TNNI2 Rabbit mAb [4H42] Images



Western blot (SDS PAGE) analysis of extracts from Human skeletal muscle tissue lyaste.using TNNI2 Rabbit mAb [4H42] at dilution of 1:1000 incubated at 4° C over

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.