

## KRTHB3 Rabbit mAb [0D2E]

Cat NO.: A34291

## Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	H,M,R	P78385	54 kda	Rabbit	IgG	100ul,200ul

	Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
The optimal dilutions should be determined by the end user  Conjugate: UnConjugate Form: Liquid sensitivity: Endogenous Purification: Protein A purification Specificity: Antibody is produced by immunizing animals with a synthetic peptide of Human KRTHB3. Storage buffer and conditions: Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer). Shipped at 4°C. Store at-20°C or -80°C. Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles. Tissue specificity: Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	WB	H,M,R	P78385	54 kda	Rabbit	IgG	100ul,200ul
Conjugate: UnConjugate Form: Liquid sensitivity: Endogenous Purification: Protein A purification Specificity: Antibody is produced by immunizing animals with a synthetic peptide of Human KRTHB3. Storage buffer and conditions: Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer). Shipped at 4°C. Store at-20°C or -80°C. Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles. Tissue specificity: Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	Applications detail:			ion			
UnConjugate Form: Liquid sensitivity: Endogenous Purification: Protein A purification Specificity: Antibody is produced by immunizing animals with a synthetic peptide of Human KRTHB3. Storage buffer and conditions: Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer). Shipped at 4°C. Store at-20°C or -80°C. Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles. Tissue specificity: Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex			The optima	al dilutions shoul	d be deter	mined by the	end user
Form: Liquid sensitivity: Endogenous Purification: Protein A purification Specificity: Antibody is produced by immunizing animals with a synthetic peptide of Human KRTHB3. Storage buffer and conditions: Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer). Shipped at 4°C. Store at-20°C or -80°C. Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles. Tissue specificity: Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	Conjugate:						
Liquid sensitivity: Endogenous Purification: Protein A purification Specificity: Antibody is produced by immunizing animals with a synthetic peptide of Human KRTHB3. Storage buffer and conditions: Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer). Shipped at 4°C. Store at-20°C or -80°C. Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles. Tissue specificity: Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	UnConjugate						
sensitivity:  Endogenous  Purification:  Protein A purification  Specificity:  Antibody is produced by immunizing animals with a synthetic peptide of Human KRTHB3.  Storage buffer and conditions:  Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer).  Shipped at 4°C. Store at-20°C or -80°C.  Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.  Tissue specificity:  Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	Form:						
Endogenous  Purification:  Protein A purification  Specificity:  Antibody is produced by immunizing animals with a synthetic peptide of Human KRTHB3.  Storage buffer and conditions:  Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer).  Shipped at 4°C. Store at-20°C or -80°C.  Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.  Tissue specificity:  Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	Liquid						
Protein A purification  Specificity:  Antibody is produced by immunizing animals with a synthetic peptide of Human KRTHB3.  Storage buffer and conditions:  Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer).  Shipped at 4°C. Store at-20°C or -80°C.  Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.  Tissue specificity:  Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	sensitivity:						
Protein A purification  Specificity:  Antibody is produced by immunizing animals with a synthetic peptide of Human KRTHB3.  Storage buffer and conditions:  Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer).  Shipped at 4°C. Store at-20°C or -80°C.  Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.  Tissue specificity:  Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	Endogenous						
Specificity:  Antibody is produced by immunizing animals with a synthetic peptide of Human KRTHB3.  Storage buffer and conditions:  Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer).  Shipped at 4°C. Store at-20°C or -80°C.  Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.  Tissue specificity:  Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	<b>Purification</b> :						
Antibody is produced by immunizing animals with a synthetic peptide of Human KRTHB3.  Storage buffer and conditions:  Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer).  Shipped at 4°C. Store at-20°C or -80°C.  Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.  Tissue specificity:  Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	Protein A purifica	tion					
Storage buffer and conditions:  Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer).  Shipped at 4°C. Store at-20°C or -80°C.  Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.  Tissue specificity:  Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	Specificity:						
Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .  Shipped at 4°C. Store at-20°C or -80°C.  Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.  Tissue specificity:  Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	Antibody is produ	iced by immuniz	ing animals with a sy	ynthetic peptide	of Human	KRTHB3.	
Shipped at 4°C. Store at-20°C or -80°C.  Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.  Tissue specificity:  Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	Storage buff	er and cond	ditions:				
Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.  Tissue specificity:  Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	Antibody store in	10 mM PBS, 0.5	mg/ml BSA, 50% gly	cerol (buffer) .			
Tissue specificity: Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	Shipped at 4°C. S	tore at-20°C or -	-80°C.				
Synthesis begins in the cortex 10-15 cell layers above the apex of the dermal papilla and ends abruptly middle of the cortex	Products are valid	d for one natura	I year of receipt. Avo	id repeated free:	ze / thaw c	ycles.	
middle of the cortex	Tissue speci	ficity:					
	Synthesis begins	in the cortex 1	0-15 cell layers abo	ve the apex of t	he dermal	papilla and	ends abruptly in the
Subcellular location:	middle of the cort	ex					
	Subcellular l	ocation:					

Function:


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry/ ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



## **Validation Data:**

## KRTHB3 Rabbit mAb [0D2E] Images



Western blot (SDS PAGE) analysis of extracts from Human skin tissue lyaste.using KRTHB3 Rabbit mAb [0D2E] at dilution of 1:1000 incubated at  $4^{\circ}$ C over night

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.