AlaRS Mouse mAb[ZL22]

Cat NO. :A35900

Information:

| [| Applications | Reactivity: | UniProt ID: | MW(kDa) | Host | Isotype | Size |
|---|--------------|-------------|-------------|---------|-------|---------|------------------|
| | WB,ICC/IF | H,M,R | P49588 | 107kDa | Mouse | lgG | 50ul 100ul,200ul |

Applications detail:

| Application | Dilution | | |
|--|-------------|--|--|
| WB | 1:1000-2000 | | |
| | | | |
| ICC/IF | 1:100 | | |
| The optimal dilutions should be determined by the end user | | | |

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human AlaRS.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Subcellular location:

Cytoplasm.

Function:

Catalyzes the attachment of alanine to tRNA(Ala) in a two-step reaction: alanine is first activated by ATP to form Ala-AMP and then transferred to the acceptor end of tRNA(Ala) (PubMed:27622773, PubMed:27911835, PubMed:28493438). Also edits incorrectly charged tRNA(Ala) via its editing domain (PubMed:27622773, PubMed:27911835, PubMed:28493438)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Validation Data:

AlaRS Mouse mAb[ZL22] Images



Western blot (SDS PAGE) analysis of extracts from HeLa cells.Using AlaRS Mouse mAb IgG [ZL22] at dilution of 1:1000 incubated at 4°C over night.

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.

For Research Use Only. Not For Use In Diagnostic Procedures.