

TAG1/TAX1 Rabbit mAb [AW73]

Cat NO. :A37410

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,ICC/IF	H,M	Q02246	135 kDa	Rabbit	IgG	50ul,100ul,200ul

Applications detail:	Application	Dilution	
	WB	1:1000-2000	
	ICC/IF	1:100,	
	The optimal dilutions should be d	be determined by the end user	

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human TAG1/TAX1.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Subcellular location:

Cell membrane, Lipid-anchor, GPI-anchor.

Function:

In conjunction with another transmembrane protein, CNTNAP2, contributes to the organization of axonal domains at nodes of Ranvier by maintaining voltage-gated potassium channels at the juxtaparanodal region. May be involved in cell adhesion..

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

TAG1/TAX1 Rabbit mAb [AW73] Images



Western blot (SDS PAGE) analysis of extracts from $\;$ Fetal brain tissue lyaste.using TAG1/TAX1 Rabbit mAb [AW73] at dilution of 1:1000 incubated at 4 $^{\circ}\mathrm{C}$ over night

View more information on http://naturebios.com