

Casein Rabbit mAb [78N7]

Cat NO. :A44832

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	н	P47710	22 kda	Rabbit	IgG	50ul,100ul,200ul

Applications detail:

Application

WB

1:1000-2000

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human Casein.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Mammary gland specific. Secreted in milk.

Subcellular location:

Secreted.

Function:

Important role in the capacity of milk to transport calcium phosphate., Casoxin D acts as opioid antagonist and has vasorelaxing activity mediated by bradykinin B1 receptors.


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

Casein Rabbit mAb [78N7] Images



Western blot (SDS PAGE) analysis of extracts from Human milk tissue lyaste.using Casein Rabbit mAb [78N7] at dilution of 1:1000 incubated at 4°C over night

View more information on http://naturebios.com