

SEC24D Mouse mAb[8B5C]

Cat NO. :A44969

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,ICC/IF	н	O94855	113kDa	Mouse	IgG	100ul,200ul

Applications detail:

Application

WB

1:1000-2000

ICC/IF

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human SEC24D.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Ubiquitously expressed, with higher amounts in placenta, pancreas, heart and liver..

Subcellular location:

 $\textbf{Cytoplasmic} \quad \textbf{vesicle}, \quad \textbf{COPII-coated} \quad \textbf{vesicle} \quad \textbf{membrane}, \textbf{Peripheral} \quad \textbf{membrane} \quad \textbf{protein}, \textbf{Cytoplasmic} \quad \textbf{side}.$

 ${\bf Endoplasmic\ reticulum\ membrane, Peripheral\ membrane\ protein, Cytoplasmic\ side.\ Cytoplasm,\ cytosol.}$

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Component of the coat protein complex II (COPII) which promotes the formation of transport vesicles from the endoplasmic reticulum (ER). The coat has two main functions, the physical deformation of the endoplasmic reticulum membrane into vesicles and the selection of cargo molecules for their transport to the Golgi complex (PubMed:17499046, PubMed:20427317, PubMed:18843296). Plays a central role in cargo selection within the COPII complex and together with SEC24C may have a different specificity compared to SEC24A and SEC24B (PubMed:17499046, PubMed:20427317, PubMed:18843296). May more specifically package GPI-anchored proteins through the cargo receptor TMED10 (PubMed:20427317). May also be specific for IxM motif-containing cargos like the SNAREs GOSR2 and STX5 (PubMed:18843296)..

Validation Data:

SEC24D Mouse mAb[8B5C] Images



Western blot (SDS PAGE) analysis of extracts from Jurkat cells.Using SEC24D Mouse mAb IgG [8B5C] at dilution of 1:1000 incubated at $4^{\circ}\mathrm{C}$ over night.

View more information on http://naturebios.com