

c-Myb (S11) Rabbit mAb [RJ6B]

Cat NO. :A51618

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,	н	P10242	72 kDa	Rabbit	IgG	100ul,200ul

Applications detail:	Application	Dilution
	WB	1:1000-2000
	IHC	1:100,
	The optimal dilutions should be determined by the end user	

	The optimal anations official be determined by the official
Conjugate:	
UnConjugate	
Form:	
Liquid	
sensitivity:	
Endogenous	
Purification:	
Protein A purification	
Specificity:	

Antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues around Ser11 of Human c-Myb .

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Subcellular location:

Nucleus.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Transcriptional activator, DNA-binding protein that specifically recognize the sequence 5'-YAAC[GT]G-3'. Plays an important role in the control of proliferation and differentiation of hematopoietic progenitor cells.

Validation Data:

c-Myb (S11) Rabbit mAb [RJ6B] Images



Western blot (SDS PAGE) analysis of extracts from Ramos cells lyastes.using c-Myb (S11) Rabbit mAb [RJ6B] at dilution of 1:1000 incubated at 4° C over night

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.