

# CCT7 Mouse mAb[IW0Y]

Cat NO. :A54370

### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	H,M	Q99832	59kDa	Mouse	IgG	50ul 100ul,200ul

Applications detail:	Application	Dilution	
	WB	1:1000-2000	
	The optimal dilutions should be	determined by the end user	

$\sim$	•		•
1''	าทแ	100	to.
$\mathbf{v}$	JIII	uga	LE:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

**Purification**:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human CCT7.

### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$ 

## Tissue specificity:

## Subcellular location:

Cytoplasm.

#### Function:

Component of the chaperonin-containing T-complex (TRiC), a molecular chaperone complex that assists the folding of proteins upon ATP hydrolysis (PubMed:25467444). The TRiC complex mediates the folding of WRAP53/TCAB1, thereby regulating telomere maintenance (PubMed:25467444). The TRiC complex plays a role in the folding of actin and tubulin (Probable)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



# **Validation Data:**

# CCT7 Mouse mAb[IW0Y] Images



Western blot (SDS PAGE) analysis of extracts from A549 cells.Using CCT7 Mouse mAb IgG [IW0Y] at dilution of 1:1000 incubated at  $4^{\circ}$ C over night.

View more information on http://naturebios.com