

# TIP47 Mouse mAb[4CG3]

Cat NO. :A58316

#### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	н	O60664	47kDa	Mouse	IgG	100ul,200ul

Applications detail:

ApplicationDilutionWB1:1000-2000IHC1:100ICC/IF1:100The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human TIP47.

### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$ 

### Tissue specificity:

### Subcellular location:

Lipid droplet. Endosome membrane, Peripheral membrane protein, Cytoplasmic side. Cytoplasm.

#### Function:

Structural component of lipid droplets, which is required for the formation and maintenance of lipid storage droplets (PubMed:34077757). Required for the transport of mannose 6-phosphate receptors (MPR) from endosomes to the trans-Golgi network (PubMed:9590177)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



## **Validation Data:**

## TIP47 Mouse mAb[4CG3] Images



Western blot (SDS PAGE) analysis of extracts from Jurkat cells. Using TIP47 Mouse mAb IgG [4CG3] at dilution of 1:1000 incubated at  $4^{\circ}\mathrm{C}$  over night.

View more information on http://naturebios.com