

# RNF7 Rabbit mAb [B3H0]

Cat NO. :A64426

#### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	H,M	Q9UBF6	13 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application Dilution

WB 1:1000-2000

IHC 1:100,

ICC/IF 1:100,

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human RNF7.

## Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$ 

## Tissue specificity:

Expressed in heart, liver, skeletal muscle and pancreas. At very low levels expressed in brain, placenta and lung..

## Subcellular location:

Cytoplasm. Nucleus.

### Function:

Probable component of the SCF (SKP1-CUL1-F-box protein) E3 ubiquitin ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins involved in cell cycle progression, signal transduction and transcription (PubMed:10851089). CRLs complexes and ARIH1 collaborate in tandem to mediate ubiquitination of target proteins, ARIH1 mediating addition of the first ubiquitin on CRLs targets (By similarity). Through the RING-type zinc finger, seems to recruit the E2 ubiquitination enzyme to the complex and brings it into close proximity to the substrate. Promotes the neddylation of CUL5 via its interaction with UBE2F. May play a role in protecting cells from apoptosis induced by redox agents..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



# **Validation Data:**

# RNF7 Rabbit mAb [B3H0] Images



Western blot (SDS PAGE) analysis of extracts from HepG2 cells lyastes.using RNF7 Rabbit mAb [B3H0] at dilution of 1:1000 incubated at  $4\,^\circ\mathrm{C}$  over night

View more information on http://naturebios.com