

14-3-3 γ Rabbit mAb [4I45]

Cat NO. :A68411

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	H,M,R	P61981	27 kDa	Rabbit	IgG	100ul,200ul

Applications detail:	Application	Dilution	
	wв	1:1000-2000	
	The optimal dilutions should be d	should be determined by the end user	

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of Human 14-3-3 $\,\gamma$

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Highly expressed in brain, skeletal muscle, and heart..

Subcellular location:

Cytoplasm.

Function:

Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

14-3-3 γ Rabbit mAb [4I45] Images



Western blot (SDS PAGE) analysis of extracts from HeLa cells.Using 14-3-3 γ Rabbit mAb [4I45] at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com