

UBE1 Mouse mAb[536M]

Cat NO. :A69656

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size	
WB,IHC,ICC/IF	H,M,R	P22314	118kda	Mouse	IgG	50ul 100ul,200ul	

Applications detail:

Application

WB 1:1000-2000

IHC 1:100

ICC/IF 1:100

The optimal dilutions should be determined by the end user

					te:		
	\sim 10		~	~	-	-	
\mathbf{u}	UH	IIU	u	а	LE	=	

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human UBE1.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Detected in erythrocytes (at protein level). Ubiquitous..

Subcellular location:

Cytoplasm. Mitochondrion. Nucleus.,[Isoform 1]: Nucleus.,[Isoform 2]: Cytoplasm.

Function:

Catalyzes the first step in ubiquitin conjugation to mark cellular proteins for degradation through the ubiquitin-proteasome system (PubMed:1606621, PubMed:1447181, PubMed:33108101). Activates ubiquitin by first adenylating its C-terminal glycine residue with ATP, and thereafter linking this residue to the side chain of a cysteine residue in E1, yielding a ubiquitin-E1 thioester and free AMP (PubMed:1447181). Essential for the formation of radiation-induced foci, timely DNA repair and for response to replication stress. Promotes the recruitment of TP53BP1 and BRCA1 at DNA damage sites (PubMed:22456334)...


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

UBE1 Mouse mAb[536M] Images



Western blot (SDS PAGE) analysis of extracts from HeLa cells.Using UBE1 Mouse mAb IgG [536M] at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com