GPX2 Rabbit mAb [874H]

Cat NO. :A73478

Information:

| Applications | Reactivity: | UniProt ID: | MW(kDa) | Host | Isotype | Size |
|--------------|-------------|-------------|---------|--------|---------|-------------|
| WB,IHC | н | P18283 | 22 kDa | Rabbit | lgG | 100ul,200ul |

Applications detail:

| Application | Dilution | | | |
|-----------------------------------|--|--|--|--|
| WB | 1:1000-2000 | | | |
| нс | 1:100, | | | |
| The optimal dilutions should be o | e optimal dilutions should be determined by the end user | | | |

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human GPX2.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Mostly in liver and gastrointestinal tract, not found in heart or kidney..

Subcellular location:

Cytoplasm, cytosol.

Function:

Could play a major role in protecting mammals from the toxicity of ingested organic hydroperoxides (PubMed:8428933). Tert-butyl hydroperoxide, cumene hydroperoxide and linoleic acid hydroperoxide but not phosphatidycholine hydroperoxide, can act as acceptors (PubMed:8428933).


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Validation Data:

GPX2 Rabbit mAb [874H] Images



Western blot (SDS PAGE) analysis of extracts from Human colon cancer tissue lyaste.using GPX2 Rabbit mAb [874H] at dilution of 1:1000 incubated at 4 °C over

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.