

IVD Mouse mAb[FT70]

Cat NO. :A74352

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	н	P26440	43kDa	Mouse	IgG	100ul,200ul

Applications detail:

Application	Dilution			
WB	1:1000-2000			
IHC	1:100			
ICC/IF	1:100			
The optimal dilutions should be determined by the end user				

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human IVD.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Subcellular location:

Mitochondrion matrix.

Function:

Catalyzes the conversion of isovaleryl-CoA/3-methylbutanoyl-CoA to 3-methylbut-2-enoyl-CoA as an intermediate step in the leucine (Leu) catabolic pathway (PubMed:7640268). To a lesser extent, is also able to catalyze the oxidation of other saturated short-chain acyl-CoA thioesters as pentanoyl-CoA, hexenoyl-CoA and butenoyl-CoA (PubMed:7640268)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

IVD Mouse mAb[FT70] Images



Western blot (SDS PAGE) analysis of extracts from HEK-293 cells. Using IVD Mouse mAb IgG [FT70] at dilution of 1:1000 incubated at 4° over night.

View more information on http://naturebios.com