

AGXT Rabbit mAb [4571]

Cat NO. :A82715

Information:

Applicat	ions	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IH	С	Н,М	P21549	40 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application

WB

1:1000-2000

IHC

The optimal dilutions should be determined by the end user

\sim		•	gate:			
	n		Oto			
\sim	<i>)</i>	uu	ıaıc	-		

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human AGXT.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Liver.

Subcellular location:

Peroxisome.

Function:

Peroxisomal aminotransferase that catalyzes the transamination of glyoxylate to glycine and contributes to the glyoxylate detoxification (PubMed:10960483, PubMed:12777626, PubMed:24055001, PubMed:23229545, PubMed:26149463). Also catalyzes the transamination between L-serine and pyruvate and contributes to gluconeogenesis from the L-serine metabolism (PubMed:10347152)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

AGXT Rabbit mAb [4571] Images



Western blot (SDS PAGE) analysis of extracts from HepG2 cells lyastes.using AGXT Rabbit mAb [4571] at dilution of 1:1000 incubated at $4\,^\circ\mathrm{C}$ over night

View more information on http://naturebios.com