

## Phospho-Gab1 (Tyr627) Rabbit mAb[Y450]

Cat NO. :A87267

## Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	н	Q13480		Rabbit	IgG	100ul,200ul
			110			

		110						
Applications detail:	Applicatio	n			Dilution			
	WB				1:1000-2000			
	The optimal dilutions should be determined by the end user							
Conjugate:								
UnConjugate								
Form:								
Liquid								
sensitivity:								
Endogenous								
Purification:								
Protein A purification								
Specificity:								
Antibody is produced by immuni	zing animals with a	synthetic pepti	ide corres	ponding to r	esidues surroundin			
Tyr627 of human Gab1								
Storage buffer and cond	litions:							
Antibody store in 10 mM PBS, 0.5r	ng/ml BSA, 50% glyce	erol (buffer) .						
Shipped at 4°C. Store at-20°C or -	80°C.							
Products are valid for one natural	year of receipt. Avoid	l repeated freez	e / thaw cy	cles.				
Tissue specificity:								
Subcellular location:								
Function:								

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry/ ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Adapter protein that plays a role in intracellular signaling cascades triggered by activated receptor-type kinases. Plays a role in FGFR1 signaling. Probably involved in signaling by the epidermal growth factor receptor (EGFR) and the insulin receptor (INSR). Involved in the MET/HGF-signaling pathway (PubMed:29408807)..

## **Validation Data:**

Phospho-Gab1 (Tyr627) Rabbit mAb[Y450] Images



Western blot (SDS PAGE) analysis of extracts from T47D cells, stimulated with heregulin...Using Phospho-Gab1 (Tyr627)Rabbit mAb IgG [Y450] at dilution of 1:1000

View more information on http://naturebios.com