BID Rabbit pAb

Cat NO. :A97119

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	н	P55957	15, 22 kDa	Rabbit	lgG	100ul,200ul

Applications detail:

Application Dilution WB 1:1000-2000 The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

peptide affinity chromatography

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of Human BID

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

[Isoform 2]: Expressed in spleen, pancreas and placenta (at protein level)...,[Isoform 3]: Expressed in lung,

pancreas and spleen (at protein level)..,[Isoform 4]: Expressed in lung and pancreas (at

Subcellular location:

Cytoplasm. Mitochondrion membrane. Mitochondrion outer membrane.

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Induces caspases and apoptosis (PubMed:14583606). Counters the protective effect of BCL2 (By similarity)..., [BH3-interacting domain death agonist p15]: Induces caspase activation and apoptosis (PubMed:15661737, PubMed:32029622). Allows the release of cytochrome c (PubMed:32029622)..., [Isoform 1]: Induces ICE-like proteases and apoptosis..., [Isoform 2]: Induces ICE-like proteases and apoptosis..., [Isoform 3]: Does not induce apoptosis..., [Isoform 4]: Induces ICE-like proteases and apoptosis...

Validation Data:

BID Rabbit pAb Images



View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.