## PEX11B Rabbit mAb[E470]

Cat NO. : A98797

### Information:

| Applications | Reactivity: | UniProt ID: | MW(kDa) | Host   | Isotype | Size             |
|--------------|-------------|-------------|---------|--------|---------|------------------|
| WB           | H,M         | O96011      | 28KDa   | Rabbit | lgG     | 50ul 100ul,200ul |

#### **Applications detail:**

# Application Dilution WB 1:1000-2000 The optimal dilutions should be determined by the end user

#### Conjugate:

UnConjugate

Form:

Liquid

#### sensitivity:

Endogenous

## **Purification**:

Protein A purification

#### Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human PEX11B.

#### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

## Subcellular location:

Peroxisome membrane, Single-pass membrane protein.

#### **Function**:

Involved in peroxisomal proliferation (PubMed:9792670). May regulate peroxisome division by recruiting the dynamin-related GTPase DNM1L to the peroxisomal membrane (PubMed:12618434). Promotes membrane protrusion and elongation on the peroxisomal surface (PubMed:20826455)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

# Validation Data:

## PEX11B Rabbit mAb[E470] Images



Western blot (SDS PAGE) analysis of extracts from U-937 cells.Using PEX11B Rabbit mAb IgG [E470] at dilution of 1:1000 incubated at 4  $^{\circ}$  over night.

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.